

# **The new IPCC Special Report « Global Warming of 1.5° C »**

**Jean-Pascal van Ypersele**

**UCLouvain (Belgium), Earth & Life Institute**


**IPCC Vice-Chair from 2008 to 2015**

**Twitter: @JPvanYpersele**

**Press conference CAN-Europe,  
Brussels, 8 October 2018**

# Global Warming of 1.5°C

An IPCC special report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty.



# Global warming of 1.5°C

*A IPCC special report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, **in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty***

Proposed outline (as adopted in October 2016; report to be finalized in 2018) :

- Summary for policy makers (max 10 pages)
- Chapters :
  - ▶ 1. Framing and context
  - ▶ 2. Mitigation pathways compatible with 1.5°C in the context of sustainable development
  - ▶ 3. Impacts of 1.5°C global warming on natural and human systems
  - ▶ 4. Strengthening and implementing the global response to the threat of climate change
  - ▶ 5. Sustainable development, poverty eradication and reducing inequalities
- Boxes (integrated case studies/regional and cross-cutting themes),
- FAQs (10 pages)

# The report in numbers

91 Authors from 40 Countries


133 Contributing authors

6000 Studies

1 113 Reviewers

42 001 Comments

# Temperature spiral



Global Mean Temperature in °C relative to 1850 – 1900

Graph: Ed Hawkins (Climate Lab Book) – Data: HadCRUT4 global temperature dataset

Animated version available on <http://openclimatedata.net/climate-spirals/temperature>

Since 1950, **extreme hot days** and **heavy precipitation** have become more common




There is evidence that anthropogenic influences, including increasing atmospheric **greenhouse gas concentrations**, have changed these extremes


# How the level of global warming affects impacts and/or risks associated with the Reasons for Concern (RFCs) and selected natural, managed and human systems

Five Reasons For Concern (RFCs) illustrate the impacts and risks of different levels of global warming for people, economies and ecosystems across sectors and regions.

## Impacts and risks associated with the Reasons for Concern (RFCs)



## Impacts and risks for selected natural, managed and human systems




Confidence level for transition: L=Low, M=Medium, H=High and VH=Very high

# How the level of global warming affects impacts and/or risks associated with the Reasons for Concern (RFCs) and selected natural, managed and human systems


Five Reasons For Concern (RFCs) illustrate the impacts and risks of different levels of global warming for people, economies and ecosystems across sectors and regions.

## Impacts and risks associated with the Reasons for Concern (RFCs)



# HALF A DEGREE OF WARMING MAKES A BIG DIFFERENCE:

EXPLAINING IPCC'S 1.5°C SPECIAL REPORT



Responsibility for content: WRI

# HALF A DEGREE OF WARMING MAKES A BIG DIFFERENCE:

EXPLAINING IPCC'S 1.5°C SPECIAL REPORT

	1.5°C	2°C	2°C IMPACTS
<b>EXTREME HEAT</b> Global population exposed to severe heat at least once every five years	 <p>14%</p>	 <p>37%</p>	<b>2.6x</b> WORSE
<b>SEA-ICE-FREE ARCTIC</b> Number of ice-free summers	 <p>AT LEAST 1 EVERY 100 YEARS</p>	 <p>AT LEAST 1 EVERY 10 YEARS</p>	<b>10x</b> WORSE
<b>SEA LEVEL RISE</b> Amount of sea level rise by 2100	 <p>0.40 METERS</p>	 <p>0.46 METERS</p>	<b>.06M</b> MORE
<b>SPECIES LOSS: VERTEBRATES</b> Vertebrates that lose at least half of their range	 <p>4%</p>	 <p>8%</p>	<b>2x</b> WORSE
<b>SPECIES LOSS: PLANTS</b> Plants that lose at least half of their range	 <p>8%</p>	 <p>16%</p>	<b>2x</b> WORSE
<b>SPECIES LOSS: INSECTS</b> Insects that lose at least half of their range	 <p>6%</p>	 <p>18%</p>	<b>3x</b> WORSE

Responsibility for content: WRI


Responsibility for content: WRI




# Cumulative emissions of CO<sub>2</sub> and future non-CO<sub>2</sub> radiative forcing determine the probability of limiting warming to 1.5°C

## a) Observed global temperature change and modeled responses to stylized anthropogenic emission and forcing pathways

Global warming relative to 1850-1900 (°C)




### b) Stylized net global CO<sub>2</sub> emission pathways Billion tonnes CO<sub>2</sub> per year (GtCO<sub>2</sub>/yr)




Faster immediate CO<sub>2</sub> emission reductions limit cumulative CO<sub>2</sub> emissions shown in panel (c).

### c) Cumulative net CO<sub>2</sub> emissions Billion tonnes CO<sub>2</sub> (GtCO<sub>2</sub>)



Maximum temperature rise is determined by cumulative net CO<sub>2</sub> emissions and net non-CO<sub>2</sub> radiative forcing due to methane, nitrous oxide, aerosols and other anthropogenic forcing agents.

### d) Non-CO<sub>2</sub> radiative forcing pathways Watts per square metre (W/m<sup>2</sup>)




## Global emissions pathway characteristics

General characteristics of the evolution of anthropogenic net emissions of CO<sub>2</sub>, and total emissions of methane, black carbon, and nitrous oxide in model pathways that limit global warming to 1.5°C with no or limited overshoot. Net emissions are defined as anthropogenic emissions reduced by anthropogenic removals. Reductions in net emissions can be achieved through different portfolios of mitigation measures illustrated in Figure SPM3B.

### Global total net CO<sub>2</sub> emissions

Billion tonnes of CO<sub>2</sub>/yr




In pathways limiting global warming to 1.5°C with **no or limited overshoot** as well as in pathways with a **high overshoot**, CO<sub>2</sub> emissions are reduced to net zero globally around 2050.

Four illustrative model pathways


### Non-CO<sub>2</sub> emissions relative to 2010

Emissions of non-CO<sub>2</sub> forcers are also reduced or limited in pathways limiting global warming to 1.5°C with **no or limited overshoot**, but they do not reach zero globally.


#### Methane emissions



#### Black carbon emissions



#### Nitrous oxide emissions



#### Timing of net zero CO<sub>2</sub>

Line widths depict the 5-95th percentile and the 25-75th percentile of scenarios

Pathways limiting global warming to 1.5°C with no or low overshoot

Pathways with high overshoot


Pathways limiting global warming below 2°C (Not shown above)

## Characteristics of four illustrative model pathways

Different mitigation strategies can achieve the net emissions reductions that would be required to follow a pathway that limit global warming to 1.5°C with no or limited overshoot. All pathways use Carbon Dioxide Removal (CDR), but the amount varies across pathways, as do the relative contributions of Bioenergy with Carbon Capture and Storage (BECCS) and removals in the Agriculture, Forestry and Other Land Use (AFOLU) sector. This has implications for the emissions and several other pathway characteristics.

### Breakdown of contributions to global net CO<sub>2</sub> emissions in four illustrative model pathways

● Fossil fuel and industry ● AFOLU ● BECCS



Global indicators	P1	P2	P3	P4	Interquartile range
Pathway classification	No or low overshoot	No or low overshoot	No or low overshoot	High overshoot	No or low overshoot
CO <sub>2</sub> emission change in 2030 (% rel to 2010)	-58	-47	-41	4	[-59, 40]
in 2050 (% rel to 2010)	-93	-95	-91	-97	[-104, 91]
Kyoto-GHG emissions* in 2030 (% rel to 2010)	-50	-49	-35	-2	[-55, 38]
in 2050 (% rel to 2010)	-82	-89	-78	-80	[-93, 61]
Final energy demand** in 2030 (% rel to 2010)	-15	-5	17	39	[-12, 7]
in 2050 (% rel to 2010)	-32	2	21	44	[-11, 22]
Renewable share in electricity in 2030 (%)	60	58	48	25	(47, 65)
in 2050 (%)	77	81	63	70	(69, 87)
Primary energy from coal in 2030 (% rel to 2010)	-78	-61	-75	-59	[-78, -59]
in 2050 (% rel to 2010)	-97	-77	-73	-97	[-95, -74]
from oil in 2030 (% rel to 2010)	-37	-13	-3	86	[-34, 3]
in 2050 (% rel to 2010)	-87	-50	-81	-32	[-78, 31]
from gas in 2030 (% rel to 2010)	-25	-20	33	37	[-26, 21]
in 2050 (% rel to 2010)	-74	-53	21	-48	[-56, 6]
from nuclear in 2030 (% rel to 2010)	59	83	98	106	(44, 102)
in 2050 (% rel to 2010)	150	98	501	468	(91, 190)
from biomass in 2030 (% rel to 2010)	-11	0	36	-1	(29, 60)
in 2050 (% rel to 2010)	-16	49	121	418	(123, 263)
from non-biomass renewables in 2030 (% rel to 2010)	430	470	315	110	(243, 438)
in 2050 (% rel to 2010)	832	1327	878	1137	(575, 1300)
Cumulative CCS until 2100 (GtCO <sub>2</sub> )	0	348	667	1218	(550, 1017)
of which BECCS (GtCO <sub>2</sub> )	0	151	414	1391	(364, 662)
Land area of bioenergy crops in 2050 (million hectare)	22	93	283	724	(151, 320)
Agricultural CH <sub>4</sub> emissions in 2030 (% rel to 2010)	-24	-48	1	14	[-30, 11]
in 2050 (% rel to 2010)	-33	-69	-23	2	[-46, 23]
Agricultural N <sub>2</sub> O emissions in 2030 (% rel to 2010)	5	-26	15	3	[-21, 4]
in 2050 (% rel to 2010)	6	-26	0	39	[-26, 1]

NOTE: Indicators have been selected to show global trends identified by the Chapter 2 assessment. National and sectoral characteristics can differ substantially from the global trends shown above.

Source: IPCC Special Report on Global Warming of 1.5°C.

\* Kyoto-gas emissions are based on SAR GWP-100

\*\* Changes in energy demand are associated with improvements in energy efficiency and behaviour change

**P1:** A scenario in which social, business, and technological innovations result in lower energy demand up to 2050 while living standards rise, especially in the global South. A down-sized energy system enables rapid decarbonisation of energy supply. Afforestation is the only CDR option considered; neither fossil fuels with CCS nor BECCS are used.

**P2:** A scenario with a broad focus on sustainability including energy intensity, human development, economic convergence and international cooperation, as well as shifts towards sustainable and healthy consumption patterns, low-carbon technology innovation, and well-managed land systems with limited societal acceptability for BECCS.

**P3:** A middle-of-the-road scenario in which societal as well as technological development follows historical patterns. Emissions reductions are mainly achieved by changing the way in which energy and products are produced, and to a lesser degree by reductions in demand.

**P4:** A resource and energy-intensive scenario in which economic growth and globalization lead to widespread adoption of greenhouse-gas intensive lifestyles, including high demand for transportation fuels and livestock products. Emissions reductions are mainly achieved through technological means, making strong use of CDR through the deployment of BECCS.

Global indicators	P1	P2	P3	P4	Interquartile range
Pathway classification	No or low overshoot	No or low overshoot	No or low overshoot	High overshoot	No or low overshoot
CO <sub>2</sub> emission change in 2030 (% rel to 2010)	-58	-47	-41	4	(-53, -40)
in 2050 (% rel to 2010)	-93	-95	-91	-97	(-104, -91)
Kyoto-GHG emissions* in 2030 (% rel to 2010)	-50	-49	-35	-2	(-55, -38)
in 2050 (% rel to 2010)	-82	-89	-78	-80	(-93, -81)
Final energy demand** in 2030 (% rel to 2010)	-15	-5	17	39	(-12, 7)
in 2050 (% rel to 2010)	-32	2	21	44	(-11, 32)
Renewable share in electricity in 2030 (%)	60	58	48	25	(47, 65)
in 2050 (%)	77	81	63	70	(69, 87)
Primary energy from coal in 2030 (% rel to 2010)	-78	-61	-75	-59	(-78, -59)
in 2050 (% rel to 2010)	-97	-77	-73	-97	(-95, -74)
from oil in 2030 (% rel to 2010)	-37	-13	-3	86	(-34, 3)
in 2050 (% rel to 2010)	-87	-50	-81	-32	(-78, -31)
from gas in 2030 (% rel to 2010)	-25	-20	33	37	(-26, 21)
in 2050 (% rel to 2010)	-74	-53	21	-48	(-56, 6)
from nuclear in 2030 (% rel to 2010)	59	83	98	106	(44, 102)
in 2050 (% rel to 2010)	150	98	501	468	(91, 190)
from biomass in 2030 (% rel to 2010)	-11	0	36	-1	(29, 80)
in 2050 (% rel to 2010)	-16	49	121	418	(123, 261)
from non-biomass renewables in 2030 (% rel to 2010)	430	470	315	110	(243, 436)
in 2050 (% rel to 2010)	832	1327	878	1137	(575, 1300)
Cumulative CCS until 2100 (GtCO <sub>2</sub> )	0	348	667	1218	(556, 1017)
of which BECCS (GtCO <sub>2</sub> )	0	151	414	1391	(364, 662)
Land area of bioenergy crops in 2050 (million hectare)	22	93	283	724	(151, 320)
Agricultural CH <sub>4</sub> emissions in 2030 (% rel to 2010)	-24	-48	1	14	(-30, -11)
in 2050 (% rel to 2010)	-33	-69	-23	2	(-46, -23)
Agricultural N <sub>2</sub> O emissions in 2030 (% rel to 2010)	5	-26	15	3	(-21, 4)
in 2050 (% rel to 2010)	6	-26	0	39	(-26, 1)


NOTE: indicators have been selected to show global trends identified by the Chapter 2 assessment. National and sectoral characteristics can differ substantially from the global trends shown above.

\* Kyoto-gas emissions are based on SAR GWP-100

\*\* Changes in energy demand are associated with improvements in energy efficiency and behaviour change

Source: IPCC Special Report on Global Warming of 1.5°C

# Comparison of global emission levels in 2025 and 2030 resulting from the implementation of the intended nationally determined contributions



UNFCCC, Aggregate effect of the intended nationally determined contributions: an update

<http://unfccc.int/resource/docs/2016/cop22/eng/02.pdf>

## Indicative linkages between mitigation options and sustainable development using SDGs (The linkages do not show costs and benefits)

Mitigation options deployed in each sector can be associated with potential positive effects (synergies) or negative effects (trade-offs) with the Sustainable Development Goals (SDGs). The degree to which this potential is realized will depend on the selected portfolio of mitigation options, mitigation policy design, and local circumstances and context. Particularly in the energy-demand sector, the potential for synergies is larger than for trade-offs. The bars group individually assessed options by level of confidence and take into account the relative strength of the assessed mitigation-SDG connections.

Length shows strength of connection




The overall size of the coloured bars depict the relative strength of synergies and trade-offs between the sectoral mitigation options and the SDGs.

Shades show level of confidence



The shades depict the level of confidence of the assessed potential for Trade-offs/Synergies.



### Length shows strength of connection




The overall size of the coloured bars depict the relative for synergies and trade-offs between the sectoral mitigation options and the SDGs.

### Shades show level of confidence



Very high Low

The shades depict the level of confidence of the assessed potential for Trade-offs/Synergies.



**Synergies: Combustion of fossil fuels, wood, and biomass also cause air pollution, which kills 7 million people per year (including 500 000 in Europe) (World Health Organization, 2018)**

**Opportunity: Addressing the causes of climate change can also improve air quality and wellbeing**

# Children are particularly sensitive to air pollution



Photo: Indiatoday.in, 6-12-2017

# Tentative and personal conclusions

---

**1.5°C matters: reducing the warming, even by tenths of a °C, can make large differences for impacts, as many of these are non-linear, that is they worsen faster with warming than the warming itself.**

**The probability of extremes (heat waves, drought, floods, extreme sea level) is significantly lower in a 1.5°C world than in a 2°C world**

**1.5°C is much safer than 2°C in terms of long-term sea-level rise associated to ice-sheet processes, particularly for low-lying regions**

# Tentative and personal conclusions

**1.5°C lower impacts will make adaptation less costly than in 2°C world, even if there is a temporary overshoot above 1.5°C**

**It is very ambitious to reduce net CO<sub>2</sub> emissions fast enough (i.e 2050) to ZERO for a 1.5°C long-term average temperature above pre-industrial objective**

**There are many possible co-benefits in fighting climate change, and they would help to achieve several SDGs**

**What is needed is the political, economic, citizen's will!**

**The slower radical changes in emission patterns take place, the more we may need uncertain or risky technologies, such as large use of carbon dioxide removal from the atmosphere (possibly at the expense of food security and biodiversity)**

**“Yes, we can!”, says the IPCC**

# To go further :

- [www.climate.be/vanyp](http://www.climate.be/vanyp) : my slides (under « conferences)
- [www.ipcc.ch](http://www.ipcc.ch) : IPCC
- [www.realclimate.org](http://www.realclimate.org) : answers to the merchants of doubt arguments
- [www.skepticalscience.com](http://www.skepticalscience.com) : same
- **Twitter: @JPvanYpersele**  
**@IPCC\_CH**